



INTERNATIONAL JOURNAL OF PHARMA PROFESSIONA'S RESEARCH



A CONCISE REVIEW ON VERSATILE MEDICINAL PLANT ACHYRANTHES ASPERA: TRADITIONAL USE, PHYTOCHEMISTRY AND PHARMACOLOGICAL ACTIVITIES

Yuvraj Tiwari¹, Nirmal Morya², Shashi Pratap Singh³, Surya Prabhakar Singh⁴
^{1,2,3,4} Pranveer Singh Institute of technology (Pharmacy) Kanpur

Keywords:

Achyranthes aspera,
 Medicinal properties,
 Pharmacological activities,
 Latjeera, chemical
 constituents, Geographical
 features

Corresponding Author-

Nirmal Morya, Assistant
 Professor, Pranveer Singh
 Institute of Technology
 (Pharmacy) Kanpur
 Email:
 nirmalmorya39@gmail.com

ABSTRACT:

Achyranthes aspera is an important medicinal plant found throughout in India. Almost every part of the plant is used in traditional systems of medicines, seeds, roots and shoots are the most important parts which are used medicinally. The present article gives updated information on its phytochemical and pharmacological properties and geographical features. The review article says that there are many phytoconstituents are present in plant having pharmacological activities like antiperiodic, diuretic, purgative, laxative, antiasthmatic, hepatoprotective, anti-allergic and various other important medicinal properties. The crushed plant is used in pneumonia and infusion of the root is used as mild astringent in constipation.

The plant reported to contain many major class of compounds like fatty acids, bisdesmosidic oleanic acid, saponins based on triterpenoid, ecdysterone, oleanic acid, triacontanol, spin sterol, dihydroxy ketones, spathulenol, alkaloids, D-glucuronic, Betaine, Achyranthine and various amino acids. This acrid, bitter plant has been used as different types of medicine and commonly used by traditional physicians for the treatment of several diseases. Though almost every of its sections are used in traditional systems of medicines, seeds, roots and shoots are the most important parts which are used medicinally and give pharmacological action. The present article provides a detail and updated information on its classification, morphology, geographical sources, phytochemical and pharmacological properties. The medicinal plant *Achyranthes aspera* L. (family: *Amaranthaceae*), which is commonly found as a weed in tropical and warm climates, is upright and has several branches.

Introduction:

Nature is the source of many medicinal plants and

many numbers of medicinal drugs are isolated from these plants and natural sources¹. In India many

biodiversity are present around 45000 species of lower and higher plants are found in India. From ancient times plants are the major source of medicines in India.² Herbal Drugs plays an Important role in all traditional systems of medicines. Herbal medicine are the mixture of several pharmacological diversity.^{4 5} Many properties of Plants are investigated like antimicrobial properties ,antiseptic properties ,wound healing properties and anti-inflammatory properties etc. are investigated by the researchers.⁶ and many bioactive compounds are present in the plants. Plants consist of many potent biochemical compounds are the components of many phytomedicines. The medicinal constituents of the plant are extracted from every part of the plantlike flowers, fruits, roots, shoots, stem, bark and leaves.⁷

According to WHO about 70-80% of world Population is based on the herbal drugs as a home remedies as a source of primary health treatment. Mostly in the case of developing countries the cost of consulting the doctor and treatment is very so the most the population rely on the treatment of herbal medicines.⁹World Health Organization has made an initiative to identify all the medicinal plants used globally and they found that more than 20000 species and more than 80% of peoples in the world are depend on herbal and traditional system of medicine for primary health care.^{10,11}

There are various chemicals which are derived from the different part of the plants and are the sources of the new drugs.¹² in the modern era the people want to sift with a herbal drugs due to side effects and hypersensitive reactions of the modern drugs.¹³ Indian folk medicines are used in prescription for many therapeutic purpose such as ulcers, snakebite, diarrhea, wounds, inflammation, scabies etc.¹⁴

A medicinal plant is defined as any plant whose one or more part of the plant are used for the therapeutic purpose or which are used for the synthesis for precursors for the direct therapeutic agents. In pharmacopoeia approx. 25% of drugs are derived from the plants and several of them are synthetic analogues based on prototype which are isolated from the plants. Infections are major cause of premature death across the world. So it is required to develop the new antimicrobial drugs which contain novel pharmacological action of mechanism. Many medicinal plants have rich source of antimicrobial

agents and used in many countries as a source of potent and powerful drugs.^{15, 16}

In the present era of drug development and discovery of new drug molecules many plant products are evaluated on the basis of their traditional uses. One of the many plants which are being evaluated for their therapeutic efficacies is *Achyranthes aspera* which is commonly known as Latjeera (Hindi) & Rough Chaff tree (English). It is an erect or procumbent, annual or perennial herb and 1- 2 m in height, often with a woody base, commonly found as a weed of waysides, on roadsides^{15, 16, 17}. Although it has many medicinal properties, it is particularly used spermicidal¹⁸, antipyretic¹⁹ & as a cardiovascular agent²⁰. The ethnic and rural people of India have preserved a significant portion of traditional knowledge on the nearby flora' therapeutic properties. This information is widely utilized to cure common illnesses and disorders and is passed down through generations through oral tradition. Chirchira has played a significant role in Indian culture and traditional medicine. It has been employed in virtually every traditional medical system, including Ayurveda, Unani, and siddha.²⁰ The tribal, rural, and indigenous inhabitants of our nation have used this plant for a variety of ailments since ancient times. Chirchira, also known as *Achyranthes aspera*, is a plant in the Amaranthaceae family that includes the species *A. canescens* R. Br., *A. argentea* Decne, *A. grand folia* Moz, *A. obovata* Peter, and *A. repea* L. Numerous names, including Chirchira (Hindi), apamarga (Sanskrit), aghedi (Gujarati), apang (Bengali), nayurivi (Tamil), and kalalat, are used in the nation (Malyalam). It is a wasteland plant that grows everywhere. It has been used as a traditional remedy for ages. It has a well-known place as a therapeutic plant in several Indian medical systems. The tribal, rural, and indigenous inhabitants of our nation have used this plant for a variety of ailments since ancient times. It holds a reputed position as medicinal plant used in several Indian medical systems. It is bitter, pungent, warming, laxative, stomachic, carminative, and good for treating vomiting, bronchitis, heart illness, piles, itchy abdominal aches, ascites, dyspepsia, dysentery, blood disorders, etc., according to Ayurveda (Bhandari, 1990; Dwivedi et. al., 2007). The plant has been referenced in texts relating to Chinese and Ayurvedic medicine. Red and white variants are

IJPPR (2023), Vol. 14, Issue 1

referenced in Ayurveda. This is known as a rough flowered stem in Sanskrit. Purgative, pungent, and digestive, it is indicated in "Nighantas" as a treatment for internal organ inflammation, piles, itch, belly enlargements, and swollen cervical glands. Ash was a key ingredient in Hindu caustic alkaline remedies. Both Indian indigenous and European physicians are familiar with the plant's diuretic qualities. Different plant components are used as ingredients in several traditional medicines, sometimes in conjunction with more potent treatments.²⁰

TAXONOMICAL CLASSIFICATION

- Species Aspera²⁰
- Genus Achyranthes²⁰
- Family Amaranthaceae²⁰
- Order Caryophyllales²⁰
- Subclass Caryophyllidae²⁰
- Kingdom Plantae²⁰
- Subkingdom Tracheobinota
- Division Mangoliophyta²⁰
- Super Division Spermatophyta
- Class Mangoliopsid²⁰



(Fig.1 Plant of *Achyranthes aspera*)

GEOGRAPHICAL FEATURES

The Plant *Achyranthes aspera* is mainly found in the tropical region of Asia, Africa, America, Australia, Ceylon, Baluchistan etc. countries. In India is mainly found on the road sides, field boundaries, waste places and forests as a weed throughout India up to the altitude of 2100m and also found in south Andaman Islands.^{21, 22}

GEOGRAPHICAL SOURCE

Achyranthes aspera, a stiff, upright annual plant, belongs to the Amaranthaceae family. The branches are 1-2 m tall, trite or completely quadrangular, striate, pubescent, and have thick leaves. The stems are angular, ribbed, and solitary or branching from the base. Different regions of the world have different names for the plant. Chaff-flower, also known as prickly chaff flower, In English, achyranth a feuilles rudes, collant, gendarme Spanish terms include molotillo, rabo de gato, rabo de chango, and rabo de raton. In Arabic, these words are known as Naeem, Noeim, Mahoot, Wazer, Hangod in the Philippines, Khare-vazhun in Persian, Atkumah in Arabic, and *Achyranthes aspera* in Latin. Sanskrit names for the term include apamargah, puthkanda, Latjira, Chirchira, Lamchichra, Sonpur, Kutri, Bengali names include Apang, Gujarati names include Safad Aghedo, telugu names include uttaraene, and Chirchitaa, Shikhari, shaikharika in Ayurvedic.²²

Plant *Achyranthes aspera* has been classified in the following order;

Kingdom	:	Plantae ²²
Division	:	Mangoliophyta
Class	:	Mangoliopsida ²²
Order	:	Caryophyllales ²²
Family	:	Amaranthaceae ²²
Genus	:	Achyranthes ²²
Species	:	Aspera ²²

The plant thrives in warmer, tropical climates. It may be found in Baluchistan, Ceylon, Australia, and America, as well as tropical Asian and African nations. It mostly grows as a weed in India along the sides of roads, on unused agricultural area, particularly in uncultivated lands, and along the edges of farmed fields.. It grows up to an altitude of 2100 m *Achyranthes aspera* is found in South Andaman Islands.^{21 22}

Latin	Achyranthes aspera
Hindi	Latjira
Sanskrit	Aghata
Punjabi	Kutri
Unani	Chirchitaa
Arabian	Atkumah
Gujarati	Safead Aghedo
Telugu	Uttaraene

TRADITIONAL USES

Achyranthes aspera is a plant that has been used traditionally to treat a variety of illnesses, including cough, asthma, and COPD. It has a variety of medicinal benefits, including laxative, purgative, diuretic, antiperiodic, and pungent qualities that are beneficial for skin rashes, boils, piles, dropsy, and oedema.^{21 22} Boiled extract of crushed plant is used in the treatment of pneumonia. In constipation the infusion of roots are used in treatment. The external application of seeds and blooming buds combined with water is employed in the treatment of skin illnesses, night blindness, reptile bites, and poisonous snake bites. When a snake bites a victim, plant roots mixed with water are administered in a paste form till the victim throws up while maintaining their consciousness. To stimulate appetite and cure a variety of gastrointestinal problems, *Achyranthes aspera* and *Smilax ovalifolia* fumes are used. It is also used to cure hemorrhoids. The leaves and seeds are used as emetic, carminatives work to reduce swelling, and they aid in digestion and phlegm expulsion.²³ for the external treatment of ulcers, the plant's ash is used. Crushed leaves were applied to sore backs to relieve strain²⁶. The plant's roots are often used to make toothbrushes. Applying a mixture made from fresh leaves might provide pain alleviation for wasp bites.²³ this plant is beneficial for rheumatism, scabies, and other skin conditions. Additionally, it has calming qualities.

FOLK REMEDIES AND TRADITIONAL USES

The herb is widely used to treat various kinds of ailments. Various traditional uses of the herb are mentioned given below-

- Whole plant is decoction boiled with water for 20-30 min is taken at night shows Diuretic in renal dropsies and general anasarca²³
- Whole plant-Decoction taken two tablespoon

three times a day-Beriberi²³

- Whole plant-Decoction boiled with water and taken twice a day-Pneumonia²³
- Whole plant-Infusion in water taken thrice a day-Bronchial infection²³
- Whole plant-Powder with lukewarm water or milk taken twice a day-Blindness in cattle and rheumatism²³
- Whole plant-Ash of plant with honey twice a day-Cough²³
- Whole plant-Juice taken thrice a day-Toothache²³
- Roots- Decoction boiled with water and taken twice a day-Pneumonia²³
- Roots- Two teaspoonful powder taken once at night-Astringent and bowel complaints²³
- Roots- Two teaspoonful powder taken once at night-Stomachic and digestive²³
- Roots- Extraction of roots taken at night-Menstrual disorders²³
- Roots- Powder taken with water daily- Leprosy²³
- Roots- Paste taken with water daily-Antifertility²³
- Roots- Infusion in water taken thrice a day- Mild astringent²³
- Roots- Ashes mixed with water –Cough²³
- Roots- Powder- taken twice a daily Bleeding in delivery²³
- Leaves Juice mixed with opium- taken twice with water Syphilitic sores²³
- Leaves Juice mixed with opium taken twice with water- Gonorrhoea²³
- Leaves Juice taken with water bed time- Bowel complaint, pile, boil, stomach, skin²³
- Leaves Decoction of powdered leaves taken twice day -Early stages of diarrhoea²³
- Leaves- Fresh leaves mixed with jaggery or black peppery and garlic and made pills - taken twice a day²³
- Seeds- Raw seeds taken with water-Expectorants²³
- Seeds- Raw seeds taken thrice a day -Brain Tonic²³
- Seeds- Raw seeds taken twice a day- Bleeding piles²³
- Flowers- Crushed flowers paste taken daily – Menorrhagia²³
- Flowers- Grounded into paste as external use Snakes and reptiles bites²³
- Fruits- Unripe fruits taken thrice daily- Respiratory disease²³

MORPHOLOGY


Achyranthes aspera (Latjeera) is an erect or

procumbent, annual or perennial herb of about 12 meter in height, often with a woody base. Stems angular, ribbed, simple or branched from the base, often with tinged purple colour, branches terete or absolutely quadrangular, striate, pubescent, leaves thick [8], $3.8 - 6.3 \times 22.5 - 4.5$ cm, ovate – elliptic or obovate – rounded, finely and softly pubescent on both sides, entire, petiolate, petiole 6 – 20 mm long, flowers greenish white, numerous in axillary or terminal spikes up to 75 cm long, seeds subcylindric, truncate at the apex, rounded at the base, reddish brown.²⁴

A 0.3–1 meter tall upright plant known as Chirchira has stiff branches that are terete or completely quadrangular, striate, and pubescent. It also has few, often thick leaves that are elliptic-ovate, petiolate, acute, and whole. Flowers are many, tiny, thick auxiliary heads or spikes that are greenish white in colour. Bracts and bracteoles are persistent and finish in a spine.²⁵ The main root is long and cylindrically thick; secondary and tertiary roots are present and seem somewhat ribbed and yellowish brown in colour. The stem is yellow brownish, upright branching, cylindrically hairy, and measures around 60 cm high. The sub-cylindrical, rounded-at-the-base, black, and brilliant seeds are truncate at the apex. The plant is found all throughout India, up to an elevation of 3000 feet.²⁶ When Prasad and Bhattacharya (1961) researched the plant pharmacognostically, they found that the stomata index, palisade ratio, vein islet number, and epidermal cell count were all on the average 6.6, 9.2, and 360. When Paliwal et al. (1960) studied the shape and growth of stomata; they discovered that the leaves were amphistomatic. Trichomes were the focus of the studies of Joshi (1931) and Dastur (1935), and the influence of various moon phases on plant development was investigated by Karnick et al. (1976).²⁷ Used in part: entire plant, including the flowers, fruits, seeds, and leaves. *Achyranthes aspera* a stiff, upright annual plant belongs to the Amaranthaceae family. Angled, ribbed, simple or branching from the base, and frequently tinted purple in colour are the characteristics of the stems (1-2 m height) absolute quadrangular or terete, striate, and pubescent and with thick leaves.²⁸

PHYTOCHEMISTRY

- There are many Phytochemicals are present in the plant the major chemical constituents are



- Other chemical constituents are present in plant are ecdysterone, long alcoholic chain, 17-Penta triacontanol, 16-hydroxyl 26 methyl heptacosan-2one, 27-cyclohexyl heptacosan-7-ol, and 36 47-dihydroxy hen-pentacontan-4one. It also contains water soluble base and betaine²⁹

CHEMICAL CONSTITUENTS OF ACHYRANTHES

- Oleanic acid from seeds³¹
- Saponins A and B³²
- Saponins C and D from unripe fruits³³
- Saponins from alcoholic extract of seeds³⁰
- AA, CHO, protein, Fe, Ca, phosphorous³⁴
- Achyranthine, N-methyl pyrrolidine –3 carboxylic acid³⁵
- Water soluble base, betaine³⁶
- Vitamin C³⁷
- Ecdysterone³⁸
- Inokosterone ecdysterone³⁸




Fig.2 Ecdysterone




Fig.3 Oleanolic acid

Fig.3 Oleanolic acid

of *Achyranthes aspera* contain oleanolic acid (0.54%).³⁴

STEM

Dihydroxy ketones-36, 37dihydroxyhenpentacontan-4-one, and Triacontanol, aliphatic alcohol, 17-pentatriacontanol, pentatriacontane, 6-pentatriacontanone, Hexatriacontane, Tritriacontane, tetracontanol-2 (C₄₀H₈₂O), 4methoxyheptatriacont-1-en-10-ol (C₃₃H₇₆O), Esitosterol and spinasterol are isolated from the shoots of the plant.³⁵

Triacontanol was also isolated along with 36, 47dihydroxyhenpentacontan-4-one 21. Two long chain compounds, extracted from the shoots, have been identified as 16-hydroxy-26-methyleheptacosan-2-one and 27-cyclohexylheptacosan-7-ol 28. In the methanol extract of the aerial parts of *Achyranthes aspera*, Kunert, 200029 reported the presence of three bisdesmosidic saponins (I-III), 20hydroxyecdysone, and quercetin-3-O-D galactoside. The compounds' structures were determined using NMR spectroscopic analysis, and their full ¹H and ¹³C assignments were obtained using 2D NMR studies. From an ethyl acetate preparation of the stem of *Achyranthes aspera*, 3Acetoxy-6 benzoyloxyapangamide was discovered by Aziz et al. in 2005. With regard to *Bacillus cereus*, the extract has modest antibacterial activity.³⁶




Fig.5 spathulenol

PHYTOCHEMISTRY

Achyranthes aspera is traditionally valued as a potent medicinal agent. Chemical constituents of various parts of the plant has been isolated and identified.

ROOT

A phytoecdysteroid called 20-Hydroxyecdysone, often known as 20E or ecdysterone, was discovered in a methanolic extract of *Achyranthes aspera* roots.³³ the ethanolic preparations of roots 22 yield an aliphatic acid called n-hexacos-14- enoic acid. Numerous other compounds, including strigmasta-5, 22-dien-3E-ol, trans-13-docasenoic acid, n-hexacosanyl ndecaniate, n-hexacos-17-enoic acid, and n-hexacos11- enoic acid, are also extracted from the root. From petroleum ether, phytosterol strigmasta-5, 22-dien-3-E-ol is extracted; benzene (75:25) elutes as a colorless crystalline mass. The Liebermann Burchard test for sterols reveals a good response. Extracts from the roots

LEAVES

The volatile oil from the leaves of *Achyranthes aspera* contains thirty different chemical components. The main component is hydroquinone (57.7%), while p-benzoquinone, Asarone, spathulenol, nerol, -ionone, and eugenol. The leaves include phenolic chemicals, alkaloids, flavonoids, saponins, tannins, and flavonoids.³⁷

SEED

Triterpenoid Saponins A and B are present in the seeds, according to phytochemical analyses. Unripe fruits have been observed to contain saponins C and D-glucose, L-rhamnose, and D-glucuronic acid are the sugars that make up its carbohydrate content (Saponins A). D-galactopyranosyl ester of D-glucuronic acid is known as saponins B. Additionally; the seeds have a water-soluble base called betaine and an alkaloid called achyranthine, 10-tricosanone, 10-octacosanone, and 4-tritriacontanone. A-L-rhamnopyranosyl (1–4) is one of three oleanolic acid

IJPPR (2023), Vol. 14, Issue 1

glycosides. (Glucopyranosyl uronic acid) -L-rhamnopyranosyl-(1-4)-(Dglucopyranosyluronic acid), -(1-3)oleanolic acid -(1→3) L- rhamnopyranosyl(14)-(-D-glucopyranosyluronic acid)-(13)oleanolic acid and -oleanolic acid-28O-D-glucopyranoside Isolated from the seeds is -28-O-D-glucopyranosyl-(1-4)-Dglucopyranoside.38,3

PHARMACOLOGICAL ACTIVITIES

- Wound Healing Activity- Ethanolic and aqueous extract of *Achyranthes aspera* consists of wound healing activity which was investigated by S.Edwin in 2008. This activity was studied in two modals excision and incision wound models.40
- Anti-allergic Activity- *Achyranthes aspera* consist of wound healing activity the petroleum ether extract of plant have anti allergic activity investigated by S.B. Datir in 2009. The steroids like B-sitosterol , ecdysone and ecdysterone are responsible for activity.41
- Cardiovascular Activity- The water soluble alkaloid which was isolated from the extract of *Achyranthes aspera* named Achyanthine have activity on the heart if Dogs and Frogs. It decrease blood pressure and Heart rate, dilated blood vessels and increase the rate of respiration.42
- Bronchoprotective Activity- The ethanolic extract of *Achyranthes aspera* shows the bronchoprotective action in TDI induced occupational asthma in rats it was studied by the scientist B.R. Goyal in 2007. The total and differential leucocytes were counted in Blood and Bronchoalveolar fluid. Liver homogenate was utilized for assessment of oxidative stress and lung histological examination was performed to investigate the inflammatory status of air way. The rats were treated by *Achyranthes aspera* did not show any abnormality.43
- Diuretic Activity- Diuretic activity was investigated by the S.S. Gupta in 1972 according to him saponins present in isolated extract of *Achyranthes aspera* shows diuretic activity in adult male rat. Achyanthine 5mg/kg is act as diuretic.34

Review Article

- Anti-depressant activity- Methanolic extract of *Achyranthes aspera* leaves shows anti- depressant action on mice and rats using forced swimming test this activity was investigated by C.C. Barua.44
- Nephroprotective activity- Methanolic extract of *Achyranthes aspera* show Nephroprotective activity against lead acetate induced nephrotoxicity in male albino rat. This activity was investigated by T.Jayakumar in 2009.45
- Anti-inflammatory Activity-Alcoholic extract of roots of *Achyranthes aspera* shows anti-inflammatory activity on wistar rats using carrageenan-induced oedema and cotton pellet granuloma test it was studied by S. Vijaya kumar on 2009.46
- The alcoholic extracts of leaves and seeds show anti-inflammatory activity in rats using carrageenan-induced paw edema method and formalin model47
- Hepatoprotective Activity- The methanolic extract of aerial part of *Achyranthes aspera* shows hepatoprotective activity on toxicity which was induced by rifampicin in albino rats this activity was reported by A.R. Bafna S.H Mishra in 2004. The extract decreases the levels of SGPT, SGOT, ALKP and total bilirubin48.
- Hypoglycemic Activity- M.S. Akhtar & J.K Iqbal in 1991 studied that the aqueous and methanolic extract of powder whole plant shows the Hypoglycemic activity it was seen by normal and diabetic induced in rats after the administration of various doses forms.49

MEDICINAL USE OF DIFFERENT PARTS OF ACHYRANTHES ASPERA

In the traditional medical systems of tropical Asian and African nations, *Achyranthes aspera* is a well-known folk remedy. The entire plant as well as its various parts, including the root, seeds, leaves, roots, flowers, and fruits, has been used for medicinal purposes.50

WHOLE PLANT:

In alloxan-induced diabetic rats, Mandar et al. (2011)¹⁹ demonstrated the effects of an ethanol extract of the whole plant on a variety of haematological (RBC, WBC count, Hb%, clotting time, O₂ carrying capacity) and biochemical (blood sugar level, lipid profile) parameters. They came to the conclusion that *Achyranthes aspera* has haematinic, hypoglycemic, and antihyperlipidemic activity that can.⁵⁰ Sheep internal parasite *Rhipicephalus microplus* larvae were resistant to the antiparasitic effects of entire plant ethyl acetate extracts (dry leaf, flower, and seed extract). *paramphistomum cervi* Male albino rats were exposed to lead acetate-induced nephrotoxicity, and the methanolic extract of the whole plant demonstrated nephroprotective action .

Ophthalmia and dysentery are both treated with the plant's juice. Uma and others, 2010 examined the antinociceptive properties of *A. aspera*'s ethanolic extract (EEAA) and sought the phytochemicals that could be behind this activity, as well as potential mechanisms of action. The whole plant's aqueous extract has hepatoprotective properties. By enhancing phagocytic activity, the Hydroalcoholic extract activates the cell-mediated immune system. In experimental rats fed a diet including sesame oil, Krishnakumari and Priya (2006) examined the antihyperlipidemic efficacy of an aqueous extract of *Achyranthes aspera*.

According to Sandhyakumari, et al., 2002 ⁴¹, an ethanolic extract of *A. aspera* induced male rats to induce reproduction. Extracts from *A. aspera* have antioxidant capabilities and may operate as primary antioxidants, free radical scavengers, or both. Hepatoprotective qualities against chemically caused hepatocellular cancer are demonstrated by the reduction in the hepatic marker.⁵¹ Using larval and adult feeding, Prasad and Pathak (2011) investigated the effects of *Achyranthes aspera* (Ash) on the reproductive fitness of *Drosophila melanogaster*. When toluene diisocyanate (TDI) was used to generate occupational asthma in wistar rats, Goyal et al. (2008) examined the bronchoprotective impact of ethanolic extract. Aside from this overall additionally, the plant is used to cure bronchial infections, blindness, rheumatism, cough, renal dropsy, beriberi, and pneumonia.⁵²

STEM/AERIAL PART

Bhattacharai, 1994 noted the rat's receptiveness to the stem bark extract's abortifacient effects. *Bacillus subtilis* and *Staphylococcus aureus* bacterial strain 49 were both inhibited by the ethanolic extract of stem. From an ethyl acetate extract of the stem of *Achyranthes aspera*, 3-Acetoxy-6-benzoyloxyapangamide was obtained by Aziz et al. in 2005. It was discovered that the extract has antibacterial properties against *Bacillus cereus*.⁵³

On rifampicin-induced hepatotoxicity in albino rats, Bafna and Mishra, 2004 ⁵⁰ reported hepatoprotective activity of the methanolic extract of the aerial parts, which reduced levels of serum glutamic pyruvic transaminase (SGPT), serum glutamic oxaloacetic transaminase (SGOT), Alkaline phosphatase (ALP), and total bilirubin According to Misra et al, 1992, the antifungal action of shoots against *Asperigillus carneus* is caused in large part by 17-pentatriacontanol. Identified substance from the plant 51's shoots' essential oil. Dry stem extracts' in vitro antibacterial activity against dental caries-causing microorganisms was investigated by Patil et al. in 2012.^{52 53}

Escherichia coli was resistant to the antibacterial effects of the stem's ethanol and methanol extract, which also revealed secondary metabolites such flavonoid and glucoside.⁵³ By using chemicals, Shendkar et al. (2012) synthesised activated carbon from the stem of *Achyranthes aspera* (by X-ray fluorescence spectroscopy). Many different uses for activated carbon include gas purification, gold purification, metal extraction, water purification, sewage treatment, air filters, and efficient catalysts.⁵⁴

LEAF

The leaves are used for the treatment of ophthalmic and other eye infections; it also has nephroprotective, post-coital antifertility, nephroprotective and immunomodulatory activities.⁵⁵ Shendkar et al, 2012 showed the presence of total thirteen amino acids in different mobile phases. Elumalai et al, 2009 ² evaluated antifungal activities from aqueous, ethanol and methanol extracts. The methanolic extract of the leaves showed antifertility activities such as abortifacient, estrogenicity, pituitary weight, ovarian hormone level and lipids profile in female rats, hypoglycaemic effect, analgesic antipyretic and

IJPPR (2023), Vol. 14, Issue 1

diuretic activities, anti-depressant effects and anti-tumor activity.⁵⁶

SEED

The leaves also include nephroprotective, post-coital antifertility, nephroprotective, and immunomodulatory properties that are used to treat ophthalmic and other eye infections. The existence of thirteen amino acids in total was demonstrated by Shendkar et al. (2012) in various mobile phases. Elumalai et al., 2009² assessed the antifungal properties of extracts in water, ethanol, and methanol.⁵⁷ The methanolic extract of the leaves demonstrated antifertility effects on female rats, including abortifacient, estrogenicity, pituitary weight, ovarian hormone level, and lipid profile, as well as hypoglycemic, analgesic, antipyretic, and diuretic effects. Antidepressant effects and tumor-fighting capacity.^{58 59}

FLOWER AND FRUIT:

Achyranthes aspera's flowers and fruits are used to treat menorrhagia, respiratory conditions, and snake and reptile bites.⁶⁰

CONCLUSION

Herbs are the natural pharmacologically active drugs used to retrieve the adjustment made in normal physiological system by foreign particles and microorganism or by any multifunctioning of the body. In every ethnic group there exists a traditional health care system, which is culturally specimen. In rural communities, health care seems to be the first and important line of defense. The WHO has already acknowledged the contribution of traditional health care in multicultural communities. It is very essential to have a proper documentation of medicinal plants and to know their potential and pharmacological activities for the improvement of health and hygiene through a traditional system of medicine.

Thus importance should be given to the potentiality of ethno medicinal studies as these can provide a very effective strategy for the discovery of useful pharmacologically active identity. A complete and systematic study is essential for identification, cataloguing and documentation of plants, which may provide a efficient and reliable way for the promotion of the traditional knowledge of the herbal medicinal plants. The present review describe that the herb

Review Article

Chirchira is used in treating several diseases. It punches on all the aspects of the herb and throws the attention to set the mind of the researchers to carry out the work for development of various formulations, which can ultimately be beneficial for the human beings as well as animals.

Acknowledgement

I want to thank Director Pharmacy Dr. A K Rai, Dean Pharmacy Dr. Pranay Wal, Head of Department Dr. Ankita Wal for providing guidance and support.

REFERENCES:

1. Cragg GM and Newman DJ. Medicinals for the millennia. *Annals of the New York Academy of Sciences* 2001; 953:3-25.
2. Elumalai EK, Chandrasekaran N, Thirumalai N, Sivakumar C, Viviyana Therasa SV and David E. *Achyranthes asperaleaf* extracts inhibited fungal growth. *International journal of pharmaceutical research* 2009; 4:1576-1579.
3. Emori TG and Gaynes RP: An overview of nosocomial infections, including the role of the microbiology laboratory. *Clinical Microbiology Reviews* 1993; 6:428-42.
4. Srivastava J, Lambert J and Vietmeyer N: Medicinal plants: An expanding role in 21 development. *World Bank Technical Paper* 1996; No. 320.
5. Uniyal SK, Singh KN, Jamwal P and Lal B: Traditional use of medicinal plants among the tribal communities of Chhota Bhangal, Western Himalaya. *Journal of Ethnobiology and Ethnomedicine* 2006; 2: 1-14.
6. Lee HS: Growth inhibitory effect of various medicinal plants against lactic acid and harmful intestinal bacteria. *Food Science and Biotechnology* 2000; 9:52-56.
7. Makari HK, Haraprasad N, Patil HS and Ravikumar: In Vitro Antioxidant Activity of the Hexane and Methanolic Extracts of *Cordia Wallichii* and *Celastrus Paniculata*. *The Internet Journal of Aesthetic and Antiaging Medicine* 2008; 1:1-10.
8. Mukherjee PK: Quality control of herbal drugs – An Approach to evaluation of Botanicals. *Business horizon pharmaceutical publishers* 2008; 13.

9. Chan K: Some aspects of toxic contaminants in herbal medicines. *Chemosphere* 2003; 52: 1371.
10. un Vijayan, Liju VB, Reena John JV, Parthipan B and Renuka C: *Indian Journal of Traditional knowledge* 2007; 6: 589-594.
11. Pandey MM, Rastogi S and Rawat AK: *The Internet Journal of Alternative Medicine* 2008; 6: 1-10.
12. Tijani Y, Uguru MO and Salawu OA: *African Journal of Biotechnology* 2008; 7: 696700.
13. Sumeet D, Raghvendra D and Kushagra M: *Achyranthesaspera Linn. (Chirchira): A magic herb in folk medicine. Ethnobotanical Leaflets* 2008; 12: 670-676.
14. Biswas TK, Maity LN and Mukherjee B: Wound healing potential of *Pterocarpus santalinus Linn*: a pharmacological evaluation. *The International Journal of Lower Extremity Wounds* 2004; 3: 143-150.
15. Jitendra B. Jain, Sheetal C. Kumane, S Bhattacharya. *Indian Journal of Traditional Knowledge*. 2006, 5(2), 237-242.
16. Anonymous. *The Wealth of India - Raw Materials*, Council of Scientific & Industrial Research, New Delhi, 2005, 55-57.
17. R. Zafar. *Medicinal Plants of India*. CBS publishers & distributors, 2009, 1-15.
18. D. Paul, D. De, K.M. Ali, K. Chatterjee, D.K. Nandi, D. Ghosh. *Contraception*, 2010, 81(4), 355-361.
19. N.G. Sutar, U.N. Sutar, Y.P. Sharma, I.K. Shaikh, S.S. Kshirsagar. *Biosciences 22 Biotechnology Research Asia*, 2008, 5(2), 841-844
20. N. C. Neogi, R. D. Garg, R. S. Rathor. *Indian Journal of Pharmacy*, 1970, 32(2), 43-46.
21. Anonymous. *The Wealth of India - Raw Materials*, Council of Scientific & Industrial Research, New Delhi, 2005, 55-57.
22. R.K. Gupta. *Medicinal & Aromatic Plants*. CBS publishers & distributors, 2010, 190.
23. R.K. Gupta. *Medicinal & Aromatic Plants*. CBS publishers & distributors, 2010, 190.
24. K.M. Nadkarni. *Indian Materia Medica*. Bombay Popular Prakashan, 2009, Vol.I, 21
25. N.K. Bhattaraj. *Fitoterapia* (1992), 63(6), 497-506
26. V.K.Singh, Z.A. Ali, S.T.H. Zaidi. *Fitoterapia* (1996), 67(2), 129-139.
27. C.P. Khare. *Indian medicinal plants*. Springer, 2007, 11-13.
28. Anonymous. *The Wealth of India - Raw Materials*, Council of Scientific & Industrial Research (CSIR), New Delhi, 2007, 17-18.
29. Dwivedi Sumeet, Dubey Raghvendra and Mehta Kushagra -review article -12sept 2008
30. Gopalanchari and Dhar –review article on traditional herbal drugs ,Kolkata(1958)
31. Khastgir et. al. *Biosciences Biotechnology Research Asia* (1950)
32. Hariharan and Rangaswami, *Pharmacognosy , Tamilnadu* (1970)
33. Sheshadri et al. *Pharmaceutical Biochemistry , SMS publishers and distributors* (1981)
34. Satyanaryana et. al. *Medicinal Plants of India*. CBS publishers & distributors (1964) 6.
35. Basu, *The Wealth of India - Raw Materials*, Council of Scientific & Industrial Research, New Delhi, 2005, 55-57.
36. Kappor and Singh, *Quality control of herbal drugs –An Approach to evaluation of Botanicals*. Business horizon pharmaceutical publishers ,(1966)
37. Hasan, *Medicinal Plants of India*. CBS publishers & distributors (1962) .
38. Chandha, *Quality control of herbal drugs –An Approach to evaluation of Botanicals*. Business horizon pharmaceutical publishers , (1970)
39. Hiroshi et. *Research article on Latjeera plant ,noval publishers* (1971)
40. S. Edwin, E. Jarald, D.L. Edwin, A. Jain, H. Kingler, K.R. Dutt, A.A. Raj. *Pharmaceutical Biology*, 2008, 46(12), 824-828.
41. S.B. Datir, A.B. Ganjare, S.A. Nirmal, S.B. Bhawar, D.K. Bharati, M.J. Patil. *Pharmacologyonline*, 2009, 921-925.
42. N. C. Neogi, R. D. Garg, R. S. Rathor. *Indian Journal of Pharmacy*, 1970, 32(2), 43-46. 23
43. B.R. Goyal, S.G. Mahajan, R.G. Mali, R.K. Goyal, A.A. Mehta. *Global Journal of Pharmacology*, 2007
44. C.C. Barua, A. Talukdar, S.A. Begum, B. Buragohain, J.D. Roy, R.S. Borah, M. Lahkar. *Pharmacologyonline*, 2009, 2, 587-594.
45. T. Jayakumar, M.P. Sridhar, T.R. Bharathprasad, M. Ilayaraja, S. Govindasamy, M.P. Balasubramanian. *Journal of Health Science*, 2009, 55(5), 701-708.
46. S.Vijaya Kumar, P. Sankar, R. Varatharajan. *Pharmaceutical Biology*, 2009, 47(10), 973975. .

47. F.A. Mehta, B.G. Patel, S.S. Pandya, K.B. Ahir, S.B. Patel. *Pharmacologyonline*, 2009, 3, 978-985.
48. A.R. Bafna, S.H. Mishra. *Ars Pharmaceutica*, 2004, 45(4), 343-351.
49. M.S. Akhtar, J. Iqbal. *Journal of Ethnopharmacology*, 1991, 31(1), 49-57.
50. Bafna SH and Mishra: *ARS Pharmaceutica* 2004; 45: 343351.
51. Misra TN, Singh RS, Pandey HS, Prasad C and Singh BP: Antifungal essential oil and a long chain alcohol from *Achyranthes aspera*. *Phytochemistry* 1992; 31: 1811-1812.
52. Patil AG and Jobanputra AH: In-vitro antimicrobial activity of *Achyranthes aspera* stem extracts against oral pathogens. *Asian Pacific Journal of Tropical Biomedicine* 2012; 1:1-4
53. Tullanithi KM, Sharmila B and Gnanendra TS: Preliminary Phytochemical analysis and Antimicrobial activity of *Achyranthes aspera*. *International Journal of Biological Technology* 2010; 1:35-38
54. Shendkar CD, Torane CD, Mundhe KS, Bhave AA and Deshpande NR: Characterization of Activated Carbon prepared from *Achyranthes aspera* Linn. by X-ray fluorescence spectroscopy (XRF). *Journal of Natural Product and Plant Resources* 2012; 2:295-97.
55. Umamaheshwari TK and Chatterjee: *International Journal of Biomedical and Pharmaceutical Sciences* 2008; 2: 70.
56. Shendkar CD, Devare SM, Ruikar AD, Lavate SM, Jawale RW and Deshpande NR: Detection of Amino acids from the Leaves of *Achyranthes aspera*. *Biomedrx International Journal* 2013; 1:511-512.
57. Shibeshi W, Makonnen E and Zerihun L: Effect of *Achyranthes aspera* L. On fetal 24 abortion, uterine and pituitary weights, serum lipids and hormones. *African Health Sciences* 2006; 6: 108-12.
58. Sutar NG, Sutar UN, Sharma YP, IShaikh IK and Kshirsagar SS: *Biosciences Biotechnology Research Asia* 2008; 5(2); 841-844.
59. Sutar N, Alok DK, Soumya MK, Priyanka G and Susri MS: Diuretic activity of *Achyranthes aspera* leaves extracts. *International research journal of pharmacy* 2012; 3:216218.
60. Barua CC, Talukdar A, Begum SA, Buragohain B, Roy JD, Borah RS and Lahkar M: Antidepressant like effects of *Achyranthes aspera* Linn in animals models of depression. *Pharmacologyonline* 2009; 2: 587-594