



INTERNATIONAL JOURNAL OF PHARMA PROFESSIONAL'S RESEARCH



DEVELOPMENT AND EVALUATION OF TOPICAL LIPOSOMAL GEL OF LEMON OIL AND ORANGE OIL FOR SYNERGISTIC EFFECT TO ENHANCE SKIN BEAUTY

Shubhrat Maheshwari¹, Aditya Singh²

¹Department of Pharmaceutics, Faculty of Pharmaceutical Sciences, Rama University Mandhana, Bithoor Road, Kanpur, Uttar Pradesh 209217

¹Department of Pharmaceutics, Faculty of Pharmacy, Integral University, Lucknow, Uttar Pradesh 206026

Keywords: Lemon oil, orange oil, Entrapped, Liposomes, Gel, Sensational

Corresponding Author-
Shubhrat Mahshwari.

E-mail:


shubhrat.maheshwari@gmail.com

Tel. +91-63999311040

ABSTRACT:

Background: In the development of cosmeceuticals for skin care, nanotechnology is critical. Essential oils are complex mixtures of numerous derivatives, including terpenoids, aliphatic chemicals, and phenol derived compounds. Due to their numerous qualities such as virucidal, bactericidal, fungicidal, local anaesthetic, anti-inflammatory, analgesic, and so on, they have been widely utilized for a long time. Essential oils are chosen over synthetically manufactured medications because they are safer for the skin. To improve the penetration of essential oils via the skin, novel nanosystems such as liposomes, niosomes, nanoemulsions, solid lipid nanoparticles, dendrimers, and nanospheres are used. These nanocarriers allow for controlled release of natural components while also improving their stability. Furthermore, nanotechnology-based skin care products have a pleasing appearance.

Graphical Abstract



Study Aim: The aim of the study is to develop lemon oil and orange oil based liposomal gel formulation for skin care

Study design: The research objective was achieved by focusing following studies. Firstly, lemon oil and orange oil is used to synthesize the liposomes. Secondly, the liposomes were converted in to liposomal gel to find an alternative skin care. Thirdly, the characteristics of were liposomal gel studied.

Purpose: To develop another alternative and herbal option for customers.

Results: The study predicament that statistical analysis multiple comparisons were made by using Unpaired Student's t-test the statistical significance was considered at $*p < 0.05$ i.e. p value summary $*P < 0.01$.

Conclusion: The sensational study indicates that lemon oil and orange oil were successfully entrapped in a liposomes to formulate gel.

Introduction:

Herbaceutical outcomes and herbal based medicaments are a magnificent source of modish therapeutic agents and for the build out of complementary and alternative medicines over traditional drug regimen [1]. Over half of the populace of globe depends on classical medicine for healthcare

aid, more than 80% of the less developed countries [2-4]. Orange oil is a markable medicinal plant which defines a wide range of biological activities, and Sensational vesicular drug carrier term was highlighted in 1991 by Gregor Cevc the aqueous core surrounded by a complex of lipid bilayer. Greek word soma clarifies the body [5, 6]. Liposomes have enomorous advantages towards biocompatibility and biodegradability as they are made from natural phospholipids similar to liposome so protect the encapsulated drug from metabolic degradation with excessive permeation of the drugs through the skin [7]. Transfersomes fetch phospholipids, surfactants, solvent, buffering agent, dyes [8, 9]. Transfersomes formulation is the process of administration of active herbaria pharmaceutical dosage form through epidermal route [10]. Cosmetics are a broad category that includes both external and internal goods. They are commonly used to give off a pleasant odour or to alter or correct the look of the body part to which they are applied. The use of cosmetic items has skyrocketed as people become more conscious about their appearance and live longer. Cosmetics are no longer restricted to simple soaps and creams; instead, more complicated goods such as anti-ageing, skin whitening, anti-acne, and soon have taken over the market. It's no surprise that the worldwide cosmetics sector is valued USD 532 billion, with forecasted growth reaching USD 716 billion by 2025 [11]. The cosmetic and skincare industries are a huge and growing part of our society. Cosmetics are defined by

the FDA as "items intended to be applied to any part of or to the human body for beautifying, cleansing, increasing attractiveness, or making a striking appearance." The term "cosmeceutical" is derived from the words "cosmetics" and "pharmaceutical medicament." Although these items have both cosmetic and biological effects on the skin, they are promoted as cosmetic or skincare products because they purport to improve the appearance of the skin [12]. Due to consumer demand to improve the appearance of their skin, hair, and nails, cosmeceutical treatments incorporate chemicals that can prevent wrinkles, UV damage, and other ailments. Nano-cosmeceuticals is defined as a cosmetic formulation that uses nanotechnology as a delivery mechanism to improve the performance of bioactive ingredients [13-17].

MATERIALS AND METHODS

Collection of Materials

Lemon oil, Orange Oil, was procured from local drug store, Lucknow, India. Soya lecithin, cholesterol, Carbopol 934, betadine was procured from Transgene Biotech, Lucknow, India.

Method

As per Table 1, Liposomal formulation was prepared. Method of Assembly operation: Soyalecthin and cholesterol was dissolved in chloroform 10ML by using individual beaker by obtained ratio respectively, Orange oil and Lemon oil was also dissolved in a separate beaker by using solvent chloroform. Taken four RBF of 1000ML loaded with respective ratio and coded with S1, S2, S3, and S4 by the 1ml addition of dissolved preparation of Orange oil and Lemon oil individually, and the RBF attached with working instrument (Rotary Vacuum Evaporator) and allowed to rotate for one hour with temperature equilibrium 60°C. When complete solvent was evaporated than a visualized thin film was succeed at the wall of RBF and the obtained thin film was hydrated by using phosphate buffer pH6.8 [18]. On the basis of observation table 1 the formulations S1 was selected

on the basis of thin film around the RBF (figure) for the gel preparation and further studied.

Table 1. Selection of formulation by the help of Rotary Vacuum Evaporator via thin film method

S.No	Cholesterol and Soyalecthin chloroform	Orange oil + Lemon oil	Obtained [Thin film formed on the surface of RBF]
S1	1:1	1:1	√
S2	2:1	1:1	×
S3	2:1	2:1	×
S4	2:2	1:1	×

Preparation of gel

In this method the 2 mg Carbopol 934 was dissolved in 50 ml of deionized water with constant stirring and heat for 30 min after heating left for cooling. When gel was stable add the prepared formulation (Figure 2) [19-20].

Drug content

The uniformity of drug content was determined using a UV spectrophotometric method. 500 mg of gel was dissolved in 50 mL of methanol. The volumetric flask was maintained for 2 hours and then thoroughly mixed in a shaker. The solution was filtered after passing through the filter paper. The drug concentration was determined using spectrophotometry with methanol as a blank [21-24].

Measurement of density

The bulk density of transferosomes was calculated by dividing its weight by its volume. The final density of a transferosomes was estimated by using an average of three specimens with a volume of $1 \times 1 \times 1 \text{ cm}^3$ [25-28].

Particle size Analysis by Malvern

The Malvern is a particle size analyzer that uses laser diffraction to measure particle sizes ranging from 0.1

IJPPR (2023), Vol. 14, Issue 2

um to 3 mm. For analysis, only a little sample (0.25g) is required, and findings can be obtained in as little as 10 minutes per sample. The fluctuation in angular scattered light intensity is measured using a laser beam passing through a dispersed sample [29-32]. (The scattering angle of small particles is modest, whereas the scattering angle of large particles is huge).

Zeta Potential Analysis by Malvern

For the measurement of zeta potential, or electrophoretic mobility, Malvern Analytical offers leading zeta potential analyzers. The Zetasizer range offers a simple, fast, and accurate approach to measure zeta potential in a number of cell types, including inexpensive disposable folded capillary cells, in both laboratory and process environments. It improve formulation stability and shelf life while reducing formulation time and expense with zeta potential measurements. One of the fundamental characteristics known to affect stability is zeta potential, which is a measure of the level of electrostatic or charge repulsion/attraction between particles [33-36].

pH of Liposomes

The pH of the formulated liposomes was checked by using buffer solution pH 6.8. Firstly 1ml of liposomes was weight and kept in 50ml of buffer for 1 hrs. Then with the help of calibrated pH meter the pH of the obtained liposomes were identified [37].

In vitro drug release study

The release studies will be carried out in 500 ml beaker containing 100 ml of phosphate buffer pH 5.8. The beaker will be assembled on a magnetic stirrer and the medium was equilibrated at $37 \pm 50^\circ\text{C}$. Dialysis membrane will be taken and one end of the membrane was sealed. After separation of non-entrapped transferosomes dispersion was filled in the dialysis membrane and other end will closed. The dialysis membrane containing the sample was suspended in the medium. Aliquots were withdrawn (1 ml) at specific intervals, filtered and the apparatus was immediately

Research Article

replenished with same quantity of fresh buffer medium pH 5.8. [38-45].

Drug entrapment efficiency Study

The transferosomes were formed and immersed in 100 ml of phosphate buffer with a pH of 7.4 overnight at 25°C . The sample leach ate was then filtered using a 0.2 mm membrane filter after being sonicated for 10 minutes. The concentration of drug in the solution was evaluated using the UV spectrophotometric (Shimadzu, Japan) method, at 228-554nm and the percentage drug entrapment was estimated by:

Entrapment Efficiency= Amount of drug entrapped / Theoretical weight of drug $\times 100$

pH of the gel

pH meter was calibrated by the help of standard buffer solution. About 1 gm of gel mixed with 100 ml of distilled water and store for 2 hours for the measurement of pH.

Microbial study

The formulated gel was inoculated on the plates of agar media. The plate was controlled by formulated transferosomes gel at 37°C for 24 hours under incubator. After duration of time period the growth rate checked by visually [45-53].

Stability study

A stability study is a set of tests used to ensure that a pharmacological product will remain stable over time. The optimum gel formulation was kept at $5^\circ\text{C} \pm 3^\circ\text{C}$ and $30^\circ\text{C} \pm 2^\circ\text{C}$ in a well-sealed stoppard glass container. For one month, data was collected by physical appearance and with digicam microscope at regular intervals.

Statistical analysis

Graph Pad Prism was used to do the statistical analysis. Multiple comparisons were made using Unpaired Student's t-test and ANOVA which was considered significance at $p < 0.05$.

RESULT

Lemon oil and Orange oil based liposomal gel was formulated by mentioned the above technique and further it was optimized on the basis on gel with hydrophilic. A table 1 shown the loading efficacy of formulation and batch S1 was selected on the basis of optimization because it has highest entrapment among them, which making it ideal for use in gel formulations. In gel preparations, carbopol 934 was frequently used as a gelling agent.

Formation of Transferosomes by Rotatory Vacuum Evaporator method

As per Table 1, the S1 selection shows the prepared formulation having a thin layer around the round bottom flask and it was hydrated by buffer with constant hand shake for 15 minute then it was observed under digicam microscope, the tiny vacuoles was visible which provide the conformation regarding the presence of transferosomes as is shown in Figure1.




Figure 1: Microscopic image of liposomes at 45x by using Digicam microscope it seems that the prepared formulations were spherical in shape. The Liposomes was formed successfully at a surface of RBF with a smooth and creamy texture and hydrated by buffer pH 6.8 and microscopically seemed spherical vesicles were formed.

Formation of Transferosomes gel

Incorporated carbopol 934 was assure the formation of curcumin based transferosomes gel which was seemed under projection microscope as is shown in Figure2.






Figure 2: The Macroscopic view of gel was thick and clear, highly viscous appearance and the Microscopic view of gel under projection microscope that shows the vesicles of drug entrapped in a spherical shape.

Fourier Transform Infrared Spectroscopy

FTIR identifies the chemical bonds in a molecule by producing an infrared absorption spectrum. Here the curcumin has a peak around 1500 cm^{-1} assigned to C=C and a peak around 1300 cm^{-1} ascribed to C-O in phenol groups present at curcumin as is shown in Figure 3.



[A]



[B]

Figure 3: FTIR of Orange oil and Lemon oil

Drug content




Figure 4: The clear supernatant solution was collected and drug concentration was evaluated using a Calibration curve of in methanol.

Density

The density of formulation at 2% w/v is 0.063 gm/cm³. According to the findings of the density study, the developed formulation has a high degree of porosity, as evidenced by its ultralow density. Additionally, the created preparation's high porosity assures that it has a high adsorption capacity for wound secretion absorption.

Particle size Analysis by Malvern




Figure 5: Liposomes appeared as spherical, well identified, unilamellar nanovesicles at peak 585.1 and the numerical value of ideal PDI ranges from 0.0 to 1 which represents perfectly uniform sample with respect to the particle size and the polymer chains are of same molecular weight, so this graph have PDI: 0.530 which is less than 1 which means the prepared formulation are uniformed with respect to particle size. Result summary, Malvern based on Laser diffraction measure the particle size distribution here, Record 223; S1 represent the liposomes and at 585.1 shows the peak with 100 % intensity, PDI: 0.530.

Zeta Potential




Figure 6: The zeta potential represents the measurement of the surface potential of suspended particles. Particles with a zeta potential greater than ± 30 mV are considered to be stable, because the repulsive force of the same charge can avoid particle aggregation. So The results of liposomes showed that the best formulation was obtained as per mentioned graph the obtained value is ±3.98 mV, which means the formulation has stability because of no aggregation between the charged particles. Result summary Malvern based on Laser diffraction measure the Zeta potential at 0.0911 mV, with 0.00125 ms/cm conductivity and zeta deviation 3.98 mV, and Record 225; S1 represent the liposomes .

pH

The pH of the liposomes was determined successfully with respect to temperature up to 45°C and it was seemed that pH was similar at different time of interval.




Figure 7: pH of liposomes at different temperature was found 5.8 which is good for topical purpose by applying student “t-test” and all the values are expressed as Mean ± SD calculated by ANOVA summary i.e. p value summary *, (significant diff. among means [P < 0.01]).

In vitro DR

In vitro DR of Liposomes was increased by the increase of time period.




Figure 8: Graph representing *In vitro* DR of liposomes by increasing the rotation time the release property was increased values are expressed as Mean ± SD calculated by ANOVA summary i.e. p value summary *, (significant diff. among means [P < 0.01]).

DEE of Leptosomes of formulation

DEE of liposomes shows that the how much amount of drug was entrapped here the result reveals that the maximum amount of drug was entrapped but less than 100 %.




Figure 9: Graph representing DEE of formulation. Values are expressed as Mean ± SD calculated by ANOVA summary i.e. p value summary *, (significant diff. among means [P < 0.01]).

pH of gel

The resultant values reveal that the pH was slightly increased but the pH values approx 5.8 which is good for skin which in not acidic nor basic.




Figure 10: Graph representing the pH was slightly increased of prepared gel at 20°C-35°C at different days of interval up to 30th days. All values are expressed as Mean ± SD calculated by ANOVA summary i.e. p value summary *, (significant diff. among means [P < 0.05]).

Microbial Study

During microbial test the distilled water, Betadine, and drug extract was compared by gel and the resultant values reveals that gel reject the null hypothesis.




Figure 11: [1] Microbial Study of gel was done by using agar media and result was achieved by using whatsmann filter paper [2] Microbial Study of gel was done by using agar media and result was achieved by using B.subtilis (gram positive bacteria), and E.coli (gram negative bacteria).




Figure 12: Graph representing the microbial study of gel in this graph the comparative study was done with each other. All values are expressed as Mean \pm SD calculated by ANOVA summary i.e. p value summary *, (significant diff. among means [P < 0.01]).

Stability study

After one month, the physical appearance of the optimized formulation S1 was found to be unchanged in all of the situations to which it was subjected. Even One month later, the medication content was confirmed to be satisfactory. The formulation was observed under digicam microscope at different time interval there was no noticeable reduction.




Figure 18: Microscopic image of gel by using Digicam microscope at different time interval for one month which shows no noticeable changes occur during one month of storage which ensures the product stability.

DISCUSSION

Aspects of Formulation

The current work of based liposomal gel for topical formulation was studied on different parameters. Firstly, the formation development which was prepared by using cholesterol and soyalecthin in chloroform that pay a significant relief on selecting a designerable development of transferosomes because they are phospholipids and they have the property of self-assembling and biodegradable in nature with biocompatibility. The drug which is the main ingredient of the formulation with recognized pharmacological actions. Figure 1 given the assurity of formulated vesicles of microscopic structure and then the pH of the liposomes was determined successfully as per Figure 7 with respect to temperature up to 45⁰C and it was seemed that pH was similar at different time of interval. The density at 2% w/v was found to be 0.063 gm/cm³. Figure 5 showed the particle size distribution by Malvern based on laser diffraction at the record 223; S1 the peak was found at 585.1 with 100 % intensity and have PDI: 0.530 which is less than 1 which means the prepared liposomes are uniformed with respect to particle size. Figure 6 showed zeta potential at 0.0911 mV, with conductivity 0.00125 ms/cm and zeta deviation 3.98 mV, formulation means particles with a zeta potential greater than \pm 30 mV are considered to be stable, because the repulsive force of the same charge can avoid particle aggregation so the results of liposomes showed that the best formulation was obtained as per mentioned graph of the mentioned value is \pm 3.98 mV, which means the formulation has stability because of no aggregation between the charged particles. Figure 8 showed the graph representing *In vitro* release of drug by increasing the rotation time the release property was increased. Figure 9 showed the drug entrapment efficiency of formulation which is less than 100%. Then step, towards the nurtured of transferosomes gel which was obtained by the addition of gelling agent carbopol and Figure 2 showed the Microscopic view of Transferosomes gel under projection microscope with small vesicles of encapsulated drug inside. Figure 10 representing the

pH was slightly increased of prepared gel at 20°C-35°C at different days of interval. Figure 3 showed the FTIR and Figure 4 showed the drug content studies by using UV Spectroscopy. Figure 11 showed the microbial study of gel which was done by using agar media and result was achieved by using whatsmann filter paper and Figure 12 showed the graph representing the microbial study of gel with the comparative study.

CONCLUSION

The present study indicates that liposomes were successfully entrapped in a formulation and. The developed formulation was evaluated for pH, particle size, drug entrapment efficiency, and percentage drug release and animal study. Summarizing all the results we can conclude that nature has provided us the best healing product than the synthetic. So need to explore herbal system for more health benefits in human subjects. They are promising carriers for because they have antioxidant properties and they can protect and stable at 4-5°C for 6 months at least. Solubility was shows in chloroform the pH of the formulated gel was found to be 5 which confirming it is suitable for topical application because in the range of 4.8 to 5.60 which is good recommended pH for the skin. All values are expressed as Mean \pm SEM calculated by ANOVA summary i.e. p value summary *, (significant diff. among them). The therapeutic potential of transfersomes gel of curcumin explores the proposed research in the management of wound healing.

ACKNOWLEDGEMENTS

The author expresses her gratitude toward Hon. Chancellor, Prof. Syed Waseem Akhtar, Integral University and Vice-Chancellor, Prof. Javed Mussarat, Integral University for providing research environment and all necessary facilities for conducting research. The authors acknowledge the office of doctoral studies and research, Integral University for critically reviewing the manuscript and providing the manuscript number for further internal communication (IU/R&D/2022-MCN0001484).

FUNDING

This research did not receive any specific grant from any funding agencies in the public, commercial, or not-for-profit sectors.

CONFLICT OF INTEREST

The authors declare no conflict of interest. The authors alone are responsible for the content and writing of this article.

REFERENCES

1. Cordell GA and Colvard MD (2012). Natural products and traditional medicine: turning on a paradigm. *Journal of Natural Products*; 75(3):514-525.
2. Farnsworth NR, Akerele O, Bingel AS, Soejarto DD and Guo Z (1985). Medicinal plants in therapy. *Bulletin of the World Health Organization*; 63(6):965-981. 7.
3. FAO (2004). Trade in medicinal plants, Economic and Social Department, Food, and Agriculture Organization of the United Nations, Rome, 2-3.
4. Nabi NG and Shrivastava M (2015). In vitro propagation of *Psoralea corylifolia* L.-an important endangered medicinal plant. *Indian Journal of Applied and Pure Biology*; 30(2):201-205.
5. Modi CD, Bharadia PD. Transfersomes: A newdominants for transdermal drug delivery. *Am J PharmTech Res* 2012; 2:71-91
6. Karande P, Mitragotri S. Enhancement of transdermal drug delivery via synergistic action of chemicals. *Biochim Biophys Acta* 2009;1788:2362-73.
7. Honeywell-Nguyen PL, Bouwstra JA. Vesicles as a tool for transdermal and dermal delivery. *Drug Discov Today Technol* 2005; 2:67-74.
8. Kombath RV, Minumula SK, Sockalingam A, Subadhra S, Parre S, Reddy TR, et al. Criticalissues related to transfersomes - Novel vesicular system. *Acta Sci Pol Technol Aliment* 2012; 11:67-82.
9. Ashish Y. Pawar, Khanderao R. Jadhav, Laxmikant H. Chaudhari Transfersome: A Novel Technique Which Improves Transdermal Permeability. *Asian Journal of Pharmaceutics* • 2016; 10(4)(Suppl):S425.
10. Kaul, S., Gulati, N., Verma, D., Mukherjee, S., & Nagaich, U. (2018). Role of nanotechnology in cosmeceuticals: a review

- of recent advances. *Journal of pharmaceuticals*, 2018.
11. Verma, D., Gulati, N., Kaul, S., Mukherjee, S., & Nagaich, U. (2018). Protein based nanostructures for drug delivery. *Journal of pharmaceuticals*, 2018.
 12. Enhanced Transdermal delivery of indinavir sulfate via transfersomes, Sheo DM, Shweta A, Vijay KT, Ram CD, Aklavya S, Ghanshyam M, *Pharmacie Globale (IJCP)* 2010; 1(06):1-7.
 13. Swarnlata Saraf, Gunjan Jeswani, Chanchal Deep Kaur, Shailendra Saraf. Development of novel herbal cosmetic cream with *Curcuma longa* extract loaded transfersomes for antiwrinkle effect. *African J Pharm Pharmacol* 2011; 5(8):1054- 1062.
 14. Woo HJ, Carraro C, Chandler D. Assembly of extended interfaces and micelles: Charge frustrated models of amphiphilic mixtures. *Faraday Discuss.* 1996; 104:183-191.
 15. Strittmatter P, Enoch HG. Formation and properties of 1000-Å diameter, single-bilayer phospholipid vesicles. *Methods Enzymol.* 1978; 52:188-193.
 16. Brannon Peppas L. Controlled Release in the Food and Cosmetics Industries in Polymeric. *Delivery Systems American Chemical Society.* 1993; 8(3):42-52.
 17. Kirby C, Gregoriadis G. Dehydration Rehydration Vesicles. A Simple Method for High Yield Drug Entrapment in Liposomes. 1984; 2(7): 979-984.
 18. Herris V. Interactive Inc. Anti-ageing medicine, vitamins, minerals and food supplements. A public opinion survey conducted for the International Longevity Center. *Journal of Anti-Aging Medicine.* 2003; 6: 83-90.
 19. Thornefeldt C. Cosmeceuticals containing herb. Facts, fiction and future. *Dermatologic Surgery.* 2005; 37: 873-80.
 20. Grove GL. Physiologic changes in older skin. *Clin Geriatr Med* 1989; 5:115-25.
 21. Gupta MA, Gupta AK. Photodamaged skin and quality of life: reason for therapy. *J Dermatolog Treat* 1996; 7:161-4.
 22. García-Varela, R.; Ramírez, O.R.F.; Serna-Saldivar, S.O.; Altamirano, J.; Cardineau, G.A. Cancer cell specific cytotoxic effect of *Rhoeo discolor* extracts and solvent fractions. *J. Ethnopharmacol.* 190, 46-58 (2016) .
 23. Shanmugam, M.; Rane, G.; Kanchi, M.; Arfuso, F.; Chinnathambi, A.; Zayed, M.; Sethi, G. The multifaceted role of curcumin in cancer prevention and treatment. *Molecules* 20, 2728-2769 (2015) .
 24. Ni, H.; Jin, W.; Zhu, T.; Wang, J.; Yuan, B.; Jiang, J.; Liang, W.; Ma, Z. Curcumin modulates TLR4/NF-κB inflammatory signaling pathway following traumatic spinal cord injury in rats. *J. Spinal Cord Med.* 38, 199- 206 (2015) .
 25. Bhatia, A.; Singh, B.; Raza, K.; Wadhwa, S.; Katare, O.P. Tamoxifen-loaded lecithin organogel (LO) for topical application: Development, optimization and characterization. *Int. J. Pharm.* 444, 47-59 (2013) .
 26. Chaudhary, H.; Kohli, K.; Amin, S.; Rathee, P.; Kumar, V. Optimization and formulation design of gels of Diclofenac and curcumin for transdermal drug delivery by Box-Behnken statistical design. *J. Pharm. Sci.* 100, 580-593 (2011) .
 27. Trelle, S.; Reichenbach, S.; Wandel, S.; Hildebrand, P.; Tschannen, B.; Villiger, P.M.; Egge, M.; Jüni, P. Cardiovascular safety of non-steroidal anti-inflammatory drugs: network meta-analysis. *Br. Med. J.* 342, c7086 (2011) .
 28. Paul, S.; Kang, S.C. Natural polyamine inhibits mouse skin inflammation and macrophage activation. *Inflamm. Res.* 62, 681-688 (2013) .
 29. Esposito, E.; Ravani, L.; Mariani, P.; Huang, N.; Boldrini, P.; Drechsler, M.; Puglia, C. Effect of nanostructured lipid vehicles on percutaneous absorption of curcumin. *Eur. J. Pharm. Biopharm.* 86, 121-132 (2012) .
 30. Wang, X.; Jiang, Y.; Wang, Y.W.; Huang, M.T.; Ho, C.T.; Huang, Q. Enhancing anti-inflammation activity of curcumin through O/W nanoemulsions. *Food Chem.* 108, 419-424 (2008) .
 31. Huang, M.-T.; Robertson, F.M.; Lysz, T.; Ferraro, T.; Wang, Z.Y.; Georgiadis, C.A.;

- Conney, A.H. Inhibitory effects of curcumin on in vitro lpx and cox activities in mouse epidermis. *Cancer Res.* 507, 338-349 (1992) .
32. Yadav, K.S.; Maan, P.; Bhatia, S.; Yadav, N.P. Exploration of anti-inflammatory activity of turmeric and onion combination on phorbol ester induced ear inflammation in mice. *Ann. Phytomed.* 3, 50-54 (2014) .
33. Nazari-Vanani, R.; Moezi, L.; Heli, H. In vivo evaluation of a self-nanoemulsifying drug delivery system for curcumin. *Biomed. Pharmacother.* 88, 715-720 (2017) .
34. Yuan, J.; Liu, R.; Ma, Y.; Zhang, Z.; Xie, Z. Curcumin attenuates airway inflammation and airway remodeling by inhibiting NF- κ B signaling and COX-2 in cigarette smoke-induced COPD Mice. *Inflammation* 41, 1804-1814 (2018).
35. Kaur R, Garg T, Das Gupta U, et al. (2014d). Preparation and characterization of spray-dried inhalable powders containing nanoaggregates for pulmonary delivery of anti-tubercular drugs. *Artif Cells Nanomed Biotechnol* 1–6.
36. Chaudhary S, Garg T, Murthy RS, et al. (2014). Recent approaches of lipid-based delivery system for lymphatic targeting via oral route. *J Drug Target* 1–12.
37. Goyal G, Garg T, Rath G, et al. (2014). Current nanotechnological strategies for an effective delivery of drugs in treatment of periodontal disease. *Crit Rev Ther Drug Carrier Syst* 31:89–119.
38. Goll DE, Bray R, Hoekstra W. (1963). Age-associated changes in muscle composition. The isolation and properties of a collagenous residue from bovine musclea. *J Food Sci* 28:503–09.
39. Parks KL, Beckman EJ. (1996). Generation of microcellular polyurethane foams via polymerization in carbon dioxide. II: foam formation and characterization. *Polym Eng Sci* 36:2417–31.
40. Pal N, Mandal S, Shiva K, Kumar B. Pharmacognostical, Phytochemical and Pharmacological Evaluation of Mallotus

- philippensis. *Journal of Drug Delivery and Therapeutics.* 2022 Sep 20;12(5):175-81.
41. Singh A, Mandal S. Ajwain (*Trachyspermum ammi* Linn): A review on Tremendous Herbal Plant with Various Pharmacological Activity. *International Journal of Recent Advances in Multidisciplinary Topics.* 2021 Jun 9;2(6):36-8.
42. Mandal S, Jaiswal V, Sagar MK, Kumar S. Formulation and evaluation of carica papaya nanoemulsion for treatment of dengue and thrombocytopenia. *Plant Arch.* 2021;21:1345-54.
43. Mandal S, Shiva K, Kumar KP, Goel S, Patel RK, Sharma S, Chaudhary R, Bhati A, Pal N, Dixit AK. Ocular drug delivery system (ODDS): Exploration the challenges and approaches to improve ODDS. *Journal of Pharmaceutical and Biological Sciences.* 2021 Jul 1;9(2):88-94.
44. Ali SA, Pathak D, Mandal S. A REVIEW OF CURRENT KNOWLEDGE ON AIRBORNE TRANSMISSION OF COVID-19 AND THEIR RELATIONSHIP WITH ENVIRONMENT. *International Journal of Pharma Professional's Research (IJPPR).* 2023;14(1):1-5.
45. Shiva K, Mandal S, Kumar S. Formulation and evaluation of topical antifungal gel of fluconazole using aloe vera gel. *Int J Sci Res Develop.* 2021;1:187-93.
46. Vishvakarma P, Mandal S, Verma A. A REVIEW ON CURRENT ASPECTS OF NUTRACEUTICALS AND DIETARY SUPPLEMENTS. *International Journal of Pharma Professional's Research (IJPPR).* 2023;14(1):78-91.
47. Ali S, Farooqui NA, Ahmad S, Salman M, Mandal S. CATHARANTHUS ROSEUS (SADABAHAR): A BRIEF STUDY ON MEDICINAL PLANT HAVING DIFFERENT PHARMACOLOGICAL ACTIVITIES. *Plant Archives.* 2021;21(2):556-9.
48. MANDAL S, JAISWAL DV, SHIVA K. A review on marketed Carica papaya leaf extract (CPL) supplements for the treatment of dengue fever with thrombocytopenia and its

- drawback. International Journal of Pharmaceutical Research. 2020 Jul;12(3).
49. Mandal S, Vishvakarma P, Verma M, Alam MS, Agrawal A, Mishra A. Solanum Nigrum Linn: An Analysis Of The Medicinal Properties Of The Plant. Journal of Pharmaceutical Negative Results. 2023 Jan 1:1595-600.
 50. Vishvakarma P, Mandal S, Pandey J, Bhatt AK, Banerjee VB, Gupta JK. An Analysis Of The Most Recent Trends In Flavoring Herbal Medicines In Today's Market. Journal of Pharmaceutical Negative Results. 2022 Dec 31:9189-98.
 51. Mandal S, Pathak D, Rajput K, Khan S, Shiva K. THROMBOPHOB-INDUCED ACUTE URTICARIA: A CASE REPORT AND DISCUSSION OF THE CASE. International Journal of Pharma Professional's Research (IJPPR). 2022;13(4):1-4.
 52. Mandal S, Shiva K, Yadav R, Sen J, Kori R. LEIOMYOSARCOMA: A CASE REPORT ON THE PREOPERATIVE DIAGNOSTIC CRITERIA. International Journal of Pharma Professional's Research (IJPPR). 2022;13(4):1-4.
 53. Mandal S, Vishvakarma P, Mandal S. Future Aspects And Applications Of Nanoemulgel Formulation For Topical Lipophilic Drug Delivery. European Journal of Molecular & Clinical Medicine.;10(01):2023.